



Research Article

Use of grape-based stain (Vinatela) on cervical cytology: A Peruvian validation study

Jeel Moya Salazar, MT, MPH^{1,2}, Victor Rojas-Zumaran, MT^{3,4}, Carlos Vegas, PhD⁴, Amalia Salafia, BSc, M.Ed^{4,5}, Hans Contreras-Pulache, MD, PhD⁶

¹Graduate School, Universidad Privada San Juan Bautista, Peru.

²Department of Biomedical Engineering, Faculty of Engineering, Universidad Tecnológica del Perú.

³Department of Pathology, Hospital Nacional Docente Madre-Niño San Bartolomé, Lima, Peru.

⁴School of Engineering in Enology and Viticulture, Universidad Privada San Juan Bautista, Ica, Perú.

⁵Faculty of Oenology and Agroindustries, Universidad Maza, Mendoza, Argentina, Peru.

⁶South America Center for Education and Research in Public Health, Universidad Norbert Wiener, Lima, Peru.



***Corresponding author:**

Jeel Moya-Salazar,
Graduate School,
Universidad Privada San Juan
Bautista, Lima, Peru.

jeel.moya@upsjb.edu.pe
jeelmoya@gmail.com

Received: 01 May 2021

Accepted: 10 July 2023

Published: 04 September 2023

DOI

10.25259/Cytojournal_19_2021

Quick Response Code:



ABSTRACT

Objectives: The Papanicolaou's (Pap's) stain is used for cervical cancer screening. It employs toxic-carcinogenic expensive reagents, which may not be easily accessible to many communities worldwide. The objective of this study was to validate the grape-based alcohol-extracted dye (Vinatela) on normal cervical samples for the Pap test.

Material and Methods: Samples of the two grape species were collected from two vineyards through the Agroindustrial Research Institute of Universidad Privada San Juan Bautista. The dye extraction from the grape species and the dye performance to stain cells were conducted in three phases: (a) direct staining with pre-fermentation wine products, (b) direct fragmentation of grapes and direct staining with shells of the grapes, and (c) alcoholic extraction of the dye. Vinatela obtained from two species (*Vitis vinifera* "Tempranillo" and "Malbec") and posterior staining of cervical samples. We conducted a double-blind validation on 30 cervical samples.

Results: The basophilic components of the cervical cells were stained. Alcoholic extraction staining protocol had a low yield. The nuclear and cytoplasmic borders, the nuclear details, and the polymorphonuclear nuclei were stained with Vinatela and could be differentiated during nuclear coloration. The initial staining protocol was 10–20 min × (mean ~12 min) staining time. We noted a slightly better staining with *V. v. Tempranillo* as compared to *V. v. Malbec* ($P = 0.045$).

Conclusion: Cervical cells stained with Vinatela stain from two grape species cultivated in the Southern of Peru, showed basophilic nuclear details.

Keywords: Cervical cytology, Grape, Papanicolaou test, Wine industry, Cervical cancer screening

INTRODUCTION

To show the cellular and tissue components, staining techniques are required that allow the differentiation of intrinsic structures, which are not detectable to the human eye. Since cellular components (such as mitochondria and nuclear histones) have different characteristics toward the use of different dyes, due to qualitative or quantitative affinities according to electrical charge, different types of dyes are used (with different auxochromic groups and mordants) for certain cellular structures.^[1] If the success of the staining depends on these characteristics, then the




Figure 2: Staining test with Vinatela dye extracted from grape species from Ica, Peru on cervical samples. (a) Protocol 1, direct coloring with pre-fermented wine (must). (b) Protocol 2, coloration test with pH change in pre-fermented wines (must). (c) Hydroalcoholic extraction of “Vinatela” dye.

stain. Cytoplasmic staining processes were performed according to hospital protocol.^[16] The quality of staining was evaluated with the staining quality index (SQI) by three cytotechnologists and two hospital pathologists with more than 20 years of experience.^[9,10,16,18] Special attention was paid to the basophilic cellular characteristics in the evaluation with the SQI (epithelial cell nuclei, polymorphonuclear cells, and cell border).

Design and statistical analysis scheme

Descriptive and inferential statistics were used to summarize the main findings. For the evaluation of staining quality, the SQI parameters previously described were used.^[18] In addition, chi-square was used to establish differences between staining protocols, staining times, and interobserver evaluations. Cohen’s Kappa was used to establish the correlation between staining, considering a value of $P < 0.05$ as statistically significant. The data analysis was carried out in the statistical analyzer IBM SPSS v22.0 (Armonk, USA) and EPIDAT-info v2.0 (Galicia, Spain) for MacOS.

RESULTS

The first staining protocol included direct staining and modification of the pre-fermented wine (must), demonstrating a range of effectiveness [Figure 3]. Of the

two grape species used in this protocol, the *V. v. Malbec* species showed better performance and stained numerous basophilic structures observed during the macroscopic controls. The use of potassium alum (20 g), hydrogen peroxide (100 μ L), and incubation (at $35 \pm 2^\circ\text{C}$) improved the performance in each stage and for both species (*V. v. Malbec* and *V. v. Tempranillo*). After identifying the direct dyeing performance of both grape species, the pH of the solution was changed, showing that the alkalinity of the species improved the affinity of the anthocyanin dyes toward the cellular structures (mainly tests 5 and 7) [Figure 2].

From these tests, we proceeded to extract the peel of both grape varieties, for subsequent analysis and testing of dyeing capacity. In [Table 1], the amounts in grams of grape peel obtained by each species are detailed. On average, 200 g of peel were extracted in 2 kg of grapes. With these shells, the staining capabilities by direct rubbing were compared [Figure 3], showing the greater capacity of the *V. v. Malbec* species. After this test, the dye was extracted, obtaining 4 ± 0.5 mL per grape species (Vinatela solution). With this alcoholic dye, the cervical cytology samples were stained for 20 min, improving the staining capacity [Table 1].

Our results showed differences between direct staining and the Vinatela protocol, the latter being the one that showed



Figure 3: Baseline characteristics of the study. (a) Sample collection sites, UPSJB vineyards, Ica, Peru. (b) Grape species included in the study (*V. v. Tempranillo* -top - and *V. v. Malbec* - below). (c) *V. v. Tempranillo* and *Malbec* grape shells. (d) Evaluation of dyeing capacity on filter paper. (e) Extraction of grape dye under Protocol 3. (f) Comparison of dyeing capacity of grape species by drag. *V. v.*: *Vitis vinifera*.

Table 1: Compounds used in the protocols for staining cervical samples with Vinatela.

S. No.	Solutions
1.	Alcohol 95°+hydrogen peroxide
2.	Hydrogen peroxide
3.	Alcohol 95°
4.	Alkaline solution
5.	Alkaline solution+hydrogen peroxide

the best affinity and dyeing capacity. [Figure 4] shows the control of nuclear staining with each protocol in cervical smears stained with Vinatela. Then, cellular staining characteristics were observed in the nuclear details of both epithelial and polymorphonuclear cells in normal/benign smears [Figure 5]. The mean of the staining procedure was 12 min.

We found better staining characteristics between *V. v. Malbec* versus *V. v. Tempranillo* ($P = 0.045$). The SQI assessment of staining established an overall SQI of 0.89 (ideal 1). Basophilic component scores showed SQI = 0.90 for chromatic pattern and nuclear staining, SQI = 0.95 for membrane continuity, SQI = 0.80 for background, and SQI = 0.89 for staining overview.

DISCUSSION

This study demonstrated the ability to stain basophilic cellular components with the grape-based stain “Vinatela” on

cervical samples from Peruvian women. Of the two evaluated species, our results showed a better performance of *V. v. Malbec*, evidenced in the cellular characteristics evaluated macroscopically and microscopically.

The strengths of this study are supported by the extraction of a stock solution of grape dye (Vinatela) under an alcoholic extraction protocol.^[15] As previously described, there is interest in the textile industry in obtaining and using natural colorants extracted from native species of grapes, corn (*Zea mays*), etc.^[19-23] The dyes extracted from these species represent a great economic contribution to the industry of different countries, producing an annual investment of 500 million dollars in Latin America.^[14] In this sense, our findings contribute to the research in the extraction of natural dyes and exceed this previous purpose using it as a cell stain of basophilic structures. Thus, it could be used in routine clinical and pathological laboratory stains (Pap stain, basophilic stain in the carbolfuchsin technique for the determination of sex chromatin, wright for blood smears, Gram for bacteria staining, etc.). This topic requires future application research.

The limitations of this study are mainly due to the nature of its design. This is the first study with an exploratory approach that has focused its methodological development on first achieving the extraction of the grape stain, and second, demonstrating its affinity toward the basophilic components of the cervical cellular structures being able to estimate their staining quality in the macro and microscopic evaluation.




Figure 4: Photomicrographs of cervical smears stained under the three staining protocols. With protocol 1 (a-d), no staining of the cellular components was obtained with *V. v. Tempranillo* (a and b), with *V. v. Malbec* slightly higher staining results were obtained (c and d). With protocol 2, improvements in staining capacity were obtained mainly in the 5:5 dilution (e and f) with *V. v. Malbec*. Protocol 3 (Vinatela) showed better cervical cell staining capacity (g-i), the staining capacity was slightly higher with *V. v. Malbec* (h and i). *V. v.*: *Vitis vinifera*.

In this sense, the Vinatela stain obtained from *V. v. Malbec* demonstrated an optimal global SQI (0.89), being comparable with the prolonged Pap modification (SQI = 0.94), with the ultrafast Pap staining (quality index = very good), with the Cytocolor (Szczezanik modification) (SQI = 0.87), with the short Pap's eosin-thiazine stain, and with the ecological stain of Pap (Eco-Pap) (SQI = 0.94).
[9,10,16,18,24,25]

On the other hand, the development of natural colorants that can be used in pathology laboratories is part of the Sustainable Development Goals (SDG) of the United Nations for 2030.^[26] Efforts for laboratories to reduce their high proportion of daily pollutants are one of the commitments that laboratories and environmental health control organizations must prioritize since tons of toxic pollutants with high carcinogenic potential are eliminated daily causing a carbon footprint^[27] and play an irreplaceable role as submarine emissaries (destroying the marine life that irreparably worsens the depressed state of the current climate).

As we proposed with the creation of the Eco-Pap,^[10] the development of laboratories specialized in natural colorants is required, which through technology transfer could reduce the impact of environmental pollution and boost the bioeconomy of the countries of the region. This progress in the study of natural dyes cannot only address the SDGs (meeting 6 of the 17 objectives), but through its development, it contributes to improving Global Environmental Health and being able to be incorporated into the routine flow work of laboratory (mainly in the Pap, which is used as the main mass secondary prevention tool) could improve cervical cancer screening (which has increased in the past 6 years)^[28] contributing to Public Health.

Vinatela initiates a new scope of action for components derived from grapes. The textile and wine industry already generates an income, being able to benefit from the possible use of grape-based stains within pathology laboratories. In this study, native grapes were used and grown in the UPSJB vineyards, highlighting the grape species from southern Peru.




Figure 5: Photomicrographs of cervical smears stained with Vinatela-based-*V. v. Malbec*. (a) Negative for intraepithelial lesion or malignancy (NILM), intermediate squamous cells with surrounding peripheral erythrocytes. (b) NILM, navicular cells. (c) NILM, metaplastic squamous cells. (d) NILM, shift in flora suggestive of bacterial vaginosis. *V. v.*: *Vitis vinifera*.

SUMMARY

We validated the Vinatela nuclear stain, from grapes grown in Peru through the alcoholic-extraction protocol, in normal cervical samples as Pap test with optimal staining quality.

COMPETING INTERESTS STATEMENT BY ALL AUTHORS

The authors declare that they have no competing interests.

AUTHORSHIP STATEMENT BY ALL AUTHORS

All authors of this article declare that we qualify for authorship as defined by ICMJE.

Each author has participated sufficiently in the work and takes public responsibility for appropriate portions of the content of this article.

Jeel Moya-Salazar planned the study design and carried the out the sampling, staining, image analysis, and alignment and drafted the manuscript. Víctor Rojas-Zumaran planned the study design and carried the out the sampling, staining, image analysis, and alignment and drafted the manuscript. Carlos Vegas planned the study design and carried the out the sampling, and drafting and alignment of the study.

Amalia Salafia and Hans Contreras-Pulache play a role in the drafting and alignment of the study. Hans Contreras-Pulache has contribution to statistical analysis.

All authors read and approved the final manuscript.

Each author acknowledges that this final version was read and approved.

ETHICS STATEMENT BY ALL AUTHORS

This study was conducted with approval from the Institutional Ethical Committee. Authors take responsibility to maintain relevant documentation in this respect.

LIST OF ABBREVIATIONS (In alphabetic order)

SQI: Staining Quality Index
 UPSJB: San Juan Bautista Private University
V. v. “Tempranillo”: *Vitis vinifera* “Tempranillo”
V. v. “Malbec”: *Vitis vinifera* “Malbec”
 Vinatela: Grape-based stain.

EDITORIAL/PEER-REVIEW STATEMENT

To ensure the integrity and highest quality of CytoJournal publications, the review process of this manuscript was

conducted under a **double-blind model** (authors are blinded for reviewers and vice versa) through the automatic online system.

FINANCIAL SUPPORT

This study was funded by the Graduate School of the San Juan Bautista Private University (No. 016-2018-VRI-UPSJB).

REFERENCES

- Culling CF. Handbook of Histopathological and Histochemical Techniques. 3rd ed. London: Butterworth; 1974.
- Papanicolaou N, Traut HF. The diagnosis value of vaginal smear in carcinoma of the uterus. Am J Obstet Gynecol 1941;42:193-205.
- Papanicolaou G. Atlas of Exfoliative Cytology. 2nd ed. Cambridge: Harvard University Press; 1954.
- Harjes U. Pap seeking new challenges. Nat Rev Cancer 2018;18:338-9.
- Catarino R, Petignat P, Dongui G, Vassilakos P. Cervical cancer screening in developing countries at a crossroad: Emerging technologies and policy choices. World J Clin Oncol 2015;6:281-90.
- Wright TC, Stoler MH, Behrens CM, Sharma A, Zhang G, Wright TL. Primary cervical cancer screening with human papillomavirus: End of study results from the ATHENA study using HPV as the first-line screening test. Gynecol Oncol 2015;136:189-97.
- de Ruiz PA, Lazcano PE, Hernández AM. Cáncer Cervicouterino. Diagnóstico, Prevención y Control. Argentina: Editorial Médica Panamericana; 2000.
- Moya-Salazar J, Rojas-Zumaran V. Analysis of the xylene problem in South American cytology laboratories. Rev Latinoam Patol Clin Med Lab 2018;65:150-8.
- Moya-Salazar J, Rojas-Zumaran V. Eco-Pap: An ecological Papanicolaou stain for sustainable cervical cancer diagnosis. Acta Cytol 2019;63:35-43.
- Moya-Salazar J, Rojas-Zumaran V. Environmental Performance of xylene, hydrochloric acid and ammonia solution during Pap stain for diagnosing cervical cancer. J Health Pollut 2016;6:58-65.
- Chantziantoniou N, Donnelly AD, Mukherjee M, Boon ME, Austin RM. Inception and development of the Papanicolaou stain method. Acta Cytol 2017;61:266-80.
- Raju K. Evolution of Pap Stain. Biomed Res Ther 2016;3:490-500.
- Jörundsson E, Lumsden JH, Jacobs RM. Rapid staining techniques in cytopathology: A review and comparison of modified protocols for hematoxylin and eosin, Papanicolaou and Romanowsky stains. Vet Clin Pathol 1999;28:100-8.
- Reyes P, Gonzalez M. Pigmentos naturales. Innova Agro 2017;5:48-53.
- Hernández V. Extracción de Antocianina a Partir de Maíz Morado (*Zea MAYS L*) para Ser Utilizado como Antioxidante y Colorante en la Industria Alimentaria. [Thesis] Lambayeque: Facultad de Ingeniería Química e Industrias Alimentarias, Universidad Nacional Pedro Ruiz Gallo; 2016.
- Moya-Salazar J, Rojas-Zumaran V. Validation of the modification of the prolonged Papanicolaou stain for the diagnosis of cervical cancer. Acta Cytol 2016;60:79-84.
- Moya-Salazar J, Rojas-Zumaran V, Torres-Martínez R, Rosas-Vargas L. Quality of Papanicolaou test smears in screening for cervical cancer, Lima, Peru. Rev Esp Patol 2016;49:7-18.
- Rojas-Zumaran VA, Moya-Salazar JJ. From the CytoColor towards ecological Pap test: Origins. Patol Rev Latinoam 2016;54:66-75.
- Manzano NP. Extracción de Antocianinas a partir de Coronta de Maíz Morado (*Zea mays L.*) Para el Aprovechamiento de Residuos Agrícolas. [Thesis]. Ambato: Facultad de Ciencia e Ingeniería en Alimentos, Universidad Técnica de Ambato; 2016.
- Huapaya FG. Influencia de los Parámetros de Extracción de Antocianinas a Partir de Cáscara de uva (*Vitis vinifera*) utilizando el Método de Taguchi. [Thesis]. Lima: Facultad de Ingeniería y Arquitectura, Universidad Peruana Unión; 2018.
- Sotomayor R. Extraction and quantification of anthocyanins from the grains of *Zea mays L.* (purple corn). Cienc Desarro 2013;16:69-74.
- Heras I, Alvis A, Arrazola G. Optimization of the Anthocyanin Extraction Process and Evaluation of the Antioxidant Capacity of Eggplant (*Solana melonera L.*). Inform Tecnol 2013;24:93-102.
- Llavona DC. Extracción y Purificación de Antocianinas a Partir de Residuos de Uva. [Thesis]. España: Universidad de Oviedo; 2000.
- Choudhary P, Sudhamani S, Pandit A, Kiri VM. Comparison of modified ultrafast Papanicolaou stain with the standard rapid Papanicolaou stain in cytology of various organs. J Cytol 2012;29:241-5.
- Walter J, Klein C, Wehrend A. Comparison of eosin-thiazin and Papanicolaou-Shorr staining for endometrial cytologies of broodmares. Tierarztl Prax Ausg G Grosstiere Nutztiere 2011;39:358-62.
- United Nations. The Sustainable Development Goals Report. New York: United Nations Publications; 2016.
- McAlister S, Barratt AL, Bell JL, McGain F. The carbon footprint of pathology testing. Med J Aust 2020;212:377-82.
- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin 2018;68:394-424.

How to cite this article: Moya-Salazar J, Rojas-Zumaran V, Vegas C, Salafia A, Contreras-Pulache H. Use of grape-based stain (Vinatela) on cervical cytology: A Peruvian validation study. CytoJournal 2023;20:30.

HTML of this article is available FREE at:
https://dx.doi.org/10.25259/Cytojournal_19_2021

The FIRST Open Access cytopathology journal

Publish in CytoJournal and RETAIN your copyright for your intellectual property

Become Cytopathology Foundation Member to get all the benefits

Annual membership fee is nominal US \$ 50 (US \$ 1000 for life)

In case of economic hardship it is free

For details visit <https://cytojournal.com/cf-member>

PubMed indexed

FREE world wide open access

Online processing with rapid turnaround time.

Real time dissemination of time-sensitive technology.

Publishes as many colored high-resolution images

Read it, cite it, bookmark it, use RSS feed, & many----



CYTOJOURNAL

www.cytojournal.com

Peer-reviewed academic cytopathology journal





NextGen CelBloking™ Kits

**Frustrated with your cell blocks?
We have a better solution!**

Nano

Nano NextGen CelBloking™

Cell block kit to process single scattered cell specimens and tissue fragments of **any** cellularity.



PATENT PENDING



Pack #1



Pack #2

Micro

Micro NextGen CelBloking™

For cellular specimens (more than 1 ml concentrated specimen with Tissuecrit more than 50%)



PATENT PENDING



Pack #1



Pack #2